

DYNAMICAL COMPLEXITY OF DISCRETE TIME REGULATORY NETWORKS

RICARDO LIMA¹ AND EDGARDO UGALDE²

¹Centre de Physique Théorique
CNRS Luminy Case 907
13288 Marseille, France

² Instituto de Física
Universidad Autónoma de San Luis Potosí
78000 San Luis Potosí, México

1. INTRODUCTION

The functioning and development of living organisms – from bacteria to humans – are controlled by genetic regulatory networks composed of interactions between DNA, RNA, proteins, and small molecules. The size and complexity of these networks make an intuitive understanding of their dynamics difficult to attain. In order to predict the behavior of regulatory systems in a systematic way, we need modeling and simulation tools with a solid foundation in mathematics and computer science.

In spite of involving a few number of genes, regulatory network usually have complex interaction graphs. Many genes are connected to each other and the incoming and outgoing interactions highly depend on the gene. This is a major difference with spatially extended systems where a usual assumption is that the incoming and outgoing interaction do not depend on the node (translation invariance). As in neural networks models, simplifying assumptions are all-to-all interactions (globally coupled systems) or random interactions. These assumptions seem not to fit exactly with the interaction graphs constructed from biological data analysis. Specific graph models and mathematical tools for the analysis of such complex networks of intermediate size are still largely missing.

Genes are segments of DNA involved in the production of proteins. The process in which a protein is produced from the information encoded in the genes is called gene expression. During this process the genes are transcribed into mRNA by enzymes called RNA polymerase, then the resulting

mRNA functions as a template for the synthesis of a protein by another enzyme, the ribosome, in a process called translation. The level of gene expression depends on the relative activity of the protein synthesis and degradation. In order to adapt the gene expression level to the requirements of the living cell, the evolution has selected complicated mechanisms regulating the production and degradation of proteins. A simple example of this, is the control of transcription by a repressor protein binding to a regulatory site on the DNA, preventing in this way the transcription of the genes. Hence, the expression level of a given gene regulates the expression level of another one, giving rise to genetic regulatory system structured by networks of interactions between genes, proteins, and molecules.

Let us illustrate the type of interactions used to define our model by means of a simple example: the regulation of the expression of the sigma factor σ^S in *Escherichia coli*, encoded by the gene *rpoS*. The factor σ^S takes its name from the fact that it plays an important role in the adaptation to a particular stress frequently encountered by bacteria: the depletion of nutriment in the environment, which leads to a considerable slowing down of cell growth called the stationary growth phase. The protein CRP, a typical repressor-activator, specifically binds the DNA at two sites close to the major promoter of *rpoS* [15]. One of these sites overlaps with the promoter, which implies that CRP and RNA polymerase cannot simultaneously bind to the DNA. As a consequence, CRP represses the transcription of *rpoS*. The example illustrates one type of regulation that is quite widespread in bacteria: a protein binding the DNA, the regulator, prevents or favors the binding of RNA polymerase to the promoter.

Genetic regulatory networks are usually modeled by systems of coupled differential equations [6], and more particularly by systems of piecewise affine differential equations like in [8]. Finite state models, better known as logical networks, are also used (see [18], or [20]). In this paper we consider a class of models of regulatory networks which may be situated in the middle of the spectrum; they present both discrete and continuous aspects. Our models consist of a network of units, whose states are quantified by a continuous real variable. The state of each unit in the network evolves according to a contractive transformation chosen from a finite collection of possible transformations. Which particular transformation is chosen at a given time step depends on the state of the neighboring units. These kind of models are directly inspired by the systems of piecewise linear differential equations mentioned before, though they do not correspond to a time discretization of those differential equations, but rather to a natural discrete time version of them. Such a discrete time model is pertinent when delays in the communications between genes justifies discrete time evolution. Besides this, the great advantage of these kind of models is that they are suitable for a detailed study from the point of view of the theory of dynamical systems. In this framework we may formulate and answer general questions about the qualitative behavior of such a system, and eventually relate those results to the experiment.

A detailed description of the asymptotic dynamics of our models would require the use of symbolic dynamics, as it has been done in [11] for systems related to ours. As a first approximation to the complete description of the dynamics we will focus on a global characteristics: the dynamical complexity. This is a well-studied notion in the framework of the theory of dynamical systems (see [10], and [2]), and it is related to the proliferation of distinguishable temporal behaviors. The main motivation of this work is to find explicit relations between the topological structure of the regulatory network, and the growth rate of the dynamical complexity. More generally we are interested in the possible constraints imposed by the topology of the underlying network and the contractive nature of the local dynamics, on the asymptotic dynamical behavior of such kind of systems. Hopefully, results on this direction will contribute to the understanding of the observed behaviors in the biological systems inspiring our models. The first steps in this direction has been fulfilled by Theorems 3.1, 4.1 and 4.2 below.

In order to illustrate our results we show some biological motivated examples. Indeed, in subsection 2.1 we discuss the self-inhibitor, whose biological prototype is presented in [1]. Then, in section we apply our results to particular families of regulatory networks. First we consider the family of circuits which models concrete biological examples. For instance, the genetic toggle switch studied in [12] corresponds to a circuit on two vertices, while the repressilator in *Escherichia coli* [9], is modeled by a circuit on three vertices. Then, in a biological inspired four vertices network studied in [21], using the results of section 4, we show that the complexity of this particular network may correspond to that of a network on two vertices.

The paper is organised as follows. In the next section we introduce the model, we give the basic definitions which we will use in the sequel, and we present some basic facts concerning the relation between complexity and asymptotic dynamics. In Section 3 we state and prove the main result of this paper, which concerns the complexity of a discrete time regulatory network satisfying a (natural but) technical condition which we call coordinatewise injectivity. In Section 4 we explore the relationship between the complexity and the structure of the network. Section 5 is devoted to some examples, and in the last section we present some concluding remarks.

Acknowledgments. During the development of this work, E. U. was invited researcher at the *Centre de Physique Théorique de Marseille*, and benefited from the financial support of the *Ministère délégué à la recherche* through the *Université de Toulon et du Var*. Both authors thank Bastien Fernandez for his remarks, comments, and suggestions throughout this work.

2. PRELIMINARIES

To each unit in interaction we associate a label in the finite set $\{1, 2, \dots, d\}$. The interaction between units is codified by an *interaction matrix* $K \in \mathcal{M}_{d \times d}([0, 1])$. The matrix entry $K_{i,j}$ specifies the strength of the action of the unit i over the unit j . The matrix K is normalized so that, for each j , $\sum_{i=1}^d K_{i,j} = 1$. The interaction has two possible modes: activation or inhibition. The interaction mode between units is codified by an *activation matrix* $s \in \mathcal{M}_{d \times d}(\{-1, 0, 1\})$, which is compatible with the interaction matrix, i. e., $s_{i,j} = 0$ if and only if $K_{i,j} = 0$. The strength of the interaction is weighted by a *contraction rate* $a \in [0, 1]$; it is directly related to the speed of degradation of the state of a noninteracting unit. At each time step $t \in \mathbb{N}$, the state of a unit $j \in \{1, 2, \dots, d\}$ is quantified by a real value $\mathbf{x}_j^t \in [0, 1]$, and it changes in time according to the transformation

$$\mathbf{x}_j^{t+1} = a\mathbf{x}_j^t + (1-a) \sum_{i=1}^d K_{i,j} H(s_{i,j}(\mathbf{x}_i^t - T_{i,j})),$$

where $H : \mathbb{R} \rightarrow \{0, 1\}$ is the Heaviside function,

$$x \mapsto H(x) := \begin{cases} 0 & \text{if } x \leq 0 \\ 1 & \text{otherwise.} \end{cases}$$

All discontinuities of the transformation are contained in the *threshold matrix* $T \in \mathcal{M}_{d \times d}([0, 1])$, which is compatible with the interaction matrix.

We let the mapping $F : [0, 1]^d \rightarrow [0, 1]^d$ defined by

$$(1) \quad F(\mathbf{x})_j := a\mathbf{x}_j^t + (1-a) \sum_{i=1}^d K_{i,j} H(s_{i,j}(\mathbf{x}_i^t - T_{i,j}))$$

induce a discrete dynamical system on the d -dimensional unit cube, a *discrete time regulatory network*. Due to discontinuities, this dynamical system may exhibit a complex behavior characterized by the existence of several attractors whose basins of attraction are intermingled in a complicated way. Often these attractors consist of a collection of disjoint periodic orbits, but there also exist attractors with infinite cardinality.

For $a = 0$, the discrete time regulatory network is equivalent to a *logical network* [13, 19]. For $a > 0$, the model is no longer equivalent to a boolean network, and can be viewed as a discrete time analogue of the a system of delay differential equations (see [3] for details). The contraction rate a plays the role of (the inverse of) a delay parameter: the smaller a , the stronger the delay is.

2.1. The self-inhibitor. In [1] it is studied an autoregulatory system which was experimentally implemented to investigate the role of negative feedback loops in the gain of stability in genetic networks. This system can be modeled by the one-dimensional regulatory network $x \mapsto F(x) := ax + (1 - a)H(T - x)$, which is semiconjugated to a rotation [4]. As we move the parameters a and T , the associated rotation number ranges over an interval, so that the attractor is either a finite set, in the case of a rational rotation number, or a Cantor set in the irrational case. Both regimes of the self-inhibited unit, periodic and quasiperiodic, share a common feature: the number of distinguishable orbits grows at most linearly in time. To be more precise, the orbits $\{x^0, x^1 = F(x^0), \dots, x^t = F(x^{t-1}), \dots\}$ and $\{y^0, y^1 = F(y^0), \dots, y^t = F(y^{t-1}), \dots\}$ are distinguishable at time $t = \tau$ if there exists $0 \leq t \leq \tau$ such that x^t and y^t lie at opposite sides of the discontinuity, i. e., $H(T - x^t) = 1 - H(T - y^t)$. Note that due to the piecewise contractive nature of the dynamics, two indistinguishable initial conditions get exponentially close as time goes to infinity.

The asymptotic dynamics of the self-inhibited unit takes place inside the interval $I := [aT, aT + (1 - a)]$, where every initial condition get trapped after a finite number of iterations. Inside the attractor there are two classes of initial conditions, corresponding to the intervals $I_0 = [T, aT + (1 - a)]$ and $I_1 := [aT, T]$. Since the map $x \mapsto F(x) := ax + (1 - a)H(T - x)$ is injective inside I , and it is an affine contraction when restricted to I_0 or to I_1 , then at most one of the intervals $F(I_0)$, $F(I_1)$, contains the discontinuity. The other one would be completely contained in one of the original intervals I_0 or I_1 . Thus, the interval whose image lies inside one of the original intervals contains initial conditions whose orbits are indistinguishable at time $t = 1$, whereas the other interval contains initial conditions which may generate up to two distinguishable orbits. Therefore, for $t = 1$ there are at most 3 distinguishable orbits represented by the same number of disjoint intervals. Those intervals are precisely the domains of continuity of the mapping $x \mapsto F^2(x)$. We may pursue this reasoning and show by induction that for each $t \in \mathbb{N}$, the maximal cardinality of a set of initial conditions in I which generate mutually distinguishable orbits at time t , is at most $t + 2$.

For the values of a and T for which the dynamics is semiconjugated to an irrational rotation, the number of distinguishable orbits may grow indefinitely. Indeed, for those values, for each $n \in \mathbb{N}$ there exists one interval $J \subset I$ where F^n is a contraction, and such that $T \in F^t(J)$. This is why for these particular choices of a and T there are $t + 1$ domains of continuity of F^t , for each $t \in \mathbb{N}$. This is precisely the case when the asymptotic dynamics is quasiperiodic, semiconjugated to an irrational rotation (see [4] for details).

In the sequel of this paper we will extend these arguments to the general case, and in this way we will obtain upper bounds for the number of distinguishable orbits.

Let us remark that the system studied in [11] is a generalization of the self-inhibitor we present here.

2.2. The complexity. In the general d -dimensional case, given a threshold matrix $T \in \mathcal{M}_{d \times d}([0, 1])$, for each coordinate $1 \leq i \leq d$ define the partition $\mathcal{P}_i$ of $[0, 1]$ so that the map

$$\mathbf{x}_i \mapsto (H(s_{i,1}(\mathbf{x}_i - T_{i,1})), H(s_{i,2}(\mathbf{x}_i - T_{i,2})), \dots, H(s_{i,d}(\mathbf{x}_i - T_{i,d})))$$

is constant restricted to each atom of $\mathcal{P}_i$. We can see that the atoms of $\mathcal{P}_i$ are semiclosed intervals with endpoints in the set $\{T_{i,j} : j = 1 \dots, d\} \cup \{0, 1\}$. The map $F : [0, 1]^d \rightarrow [0, 1]^d$ defined by Eq. (1) is an affine contraction restricted to the atoms of the Cartesian product $\mathcal{P} := \prod_{i=1}^d \mathcal{P}_i$, which we call *base partition*. Indeed, if $\mathbf{x}, \mathbf{y} \in [0, 1]^d$ belong to the same atom of $\mathcal{P}$, then for each $1 \leq i, j \leq d$ we have $H(s_{i,j}(\mathbf{x}_i - T_{i,j})) = H(s_{i,j}(\mathbf{y}_i - T_{i,j}))$, which implies $F(\mathbf{x}) - F(\mathbf{y}) = a(\mathbf{x} - \mathbf{y})$.

We define recursively the dynamical partitions

$$(2) \quad \mathcal{P}^{t+1} := \{F^{-1}(\mathbf{I}) \cap \mathbf{J} : \mathbf{I} \in \mathcal{P}^t \text{ and } \mathbf{J} \in \mathcal{P}\},$$

with $\mathcal{P}^1 := \mathcal{P}$. It is easy to verify that the τ -th iterate of the map F , $F^\tau : [0, 1]^d \rightarrow [0, 1]^d$, is an affine contraction in each atom of $\mathcal{P}^\tau$. It means that two initial conditions in the same atom of $\mathcal{P}^\tau$ will get closer during the first τ time steps.

The following discussion holds in the slightly more general case when the mapping $F : [0, 1]^d \rightarrow [0, 1]^d$ is a non necessarily affine contraction when restricted to each atom of the base partition $\mathcal{P}$. We call this a *piecewise contraction*.

The *complexity* of the system $([0, 1]^d, F)$ is the integer valued function

$$C : \mathbb{N} \rightarrow \mathbb{N} \text{ such that } C(t) := \#\mathcal{P}^t.$$

The number of atoms in the dynamical partition $\mathcal{P}^t$ equals the number of distinguishable orbits up to time t . It is the aim of this paper to give upper bounds on the growth of this quantity. For systems defined by a continuous transformation acting on a compact metric space, if the complexity is eventually constant then all orbits are eventually periodic. For the same kind of system, to a fast growth of the complexity one can associate some notion of chaos (see [2] and [10]). The computation of the complexity is part of program whose aim is the complete characterization of the asymptotic dynamics of the discrete time regulatory network. Information about the growth of the complexity, together with a description of the recurrence properties of individual orbits, will allow us to give a detailed picture of the asymptotic behavior of the system.

2.3. The asymptotic behavior. The *limit set* Λ of the system $([0, 1]^d, F)$ is the collection of all accumulation points of its orbits, i. e., $\mathbf{x} \in \Lambda$ whenever there exists an initial condition $\mathbf{y} \in [0, 1]^d$, and a sequence of times $t_1 < t_2 < \dots < t_n < \dots$, such that $\lim_{n \rightarrow \infty} F^{(t_n)}(\mathbf{y}) = \mathbf{x}$.

Let us denote

$$\Delta := \left\{ \mathbf{x} \in [0, 1]^d : \exists i, k \in \{1, 2, \dots, d\} \text{ such that } \mathbf{x}_k = T_{i,k} \right\},$$

the discontinuity set. When $\Lambda \cap \Delta = \emptyset$, then $\Lambda := \bigcap_{t=0}^{\infty} F^t([0,1]^d)$, and it contains all the orbits with infinite past.

If the limit set has infinite cardinality, then the complexity is a strictly increasing function. Indeed, if on the contrary $C(\tau + 1) = C(\tau)$ for some $\tau \in \mathbb{N}$, then necessarily to each atom $\mathbf{I} \in \mathcal{P}^\tau$ there corresponds a unique atom $\bar{\mathbf{I}}$ such that $F(\mathbf{I}) \subset \bar{\mathbf{I}}$. The transformation $\mathbf{I} \mapsto \bar{F}(\mathbf{I}) := \bar{\mathbf{I}}$, acting over $\mathcal{P}^\tau$, would generate in this case up to $C(\tau)$ ultimately periodic orbits. Thus, to each atom $\mathbf{I} \in \mathcal{P}^\tau$ we could associate a transient $t \in \mathbb{N}$ and a period $p \in \mathbb{N}$ such that $\bar{F}^{t+kp}(\mathbf{I}) = \bar{F}^t(\mathbf{I})$ for each $k \in \mathbb{N}$. Since F is a contraction when restricted to each atom of $\mathcal{P}^\tau$, all the orbits with initial condition $\mathbf{x} \in \mathbf{I}$ would have one and the same accumulation point in the closure of each one of the atoms $\bar{F}^t(\mathbf{I}), \bar{F}^{t+1}(\mathbf{I}), \dots, \bar{F}^{t+p-1}(\mathbf{I})$. Therefore, $C(\tau + 1) = C(\tau)$ implies $\#\Lambda < \infty$. Of course, if Λ is a finite set, then the dynamical complexity has to be a bounded function.

One would expect bounded dynamical complexity to be a generic situation in the case of piecewise contractions. The following suggests this is true.

Proposition 2.1 (Multiperiodicity). *If $\text{dist}(\Lambda, \Delta) > 0$ then $\#\Lambda < \infty$.*

Proof. Let $\epsilon = \text{dist}(\Lambda, \Delta)/2 > 0$. Compactness allows us to choose a finite cover $\mathcal{C} = \{B(\mathbf{x}^{(i)}, \epsilon_i) : \mathbf{x}^{(i)} \in \Lambda, i = 1, 2, \dots, N\}$ for $\bar{\Lambda}$, composed by open balls of radius not greater than ϵ and centered at points in Λ . Since $\epsilon_i < \text{dist}(\Lambda, \Delta)$ for each $i = 1, 2, \dots, N$, then $F^t(B(\mathbf{x}^{(i)}, \epsilon_i))$ is a ball of radius $a^t \epsilon_i$ centered at $F^t(\mathbf{x}^{(i)})$, for each $t \in \mathbb{N}$. In this way, for each $t \in \mathbb{N}$, the collection $\mathcal{C}_t := \{B(F^t(\mathbf{x}^{(i)}), a^t \epsilon_i) : i = 1, 2, \dots, N\}$ is an open cover for $F^t(\bar{\Lambda})$, but since Λ is F -invariant, then $\mathcal{C}_t$ is an open cover for Λ .

Let t be such that $a^t \epsilon$ is less than or equal to the Lebesgue number of $\mathcal{C}$. Then, for each $i = 1, 2, \dots, N$ there exists $j \in \{1, 2, \dots, N\}$ such that

$$B(F^t(\mathbf{x}^{(i)}), a^t \epsilon_i) \subset B(\mathbf{x}^{(j)}, \epsilon_j).$$

For each $i = 1, 2, \dots, N$ choose one such j , and define a map $i \mapsto \tau(i) := j$ from $\{1, 2, \dots, N\}$ to itself. The successive iterations of this map, $i \mapsto \tau(i) \mapsto \dots$, generate ultimately periodic sequences, but the F -invariance of Λ implies that each $i \in \{1, 2, \dots, N\}$ has to be periodic under τ . This means that for each $i \in \{1, 2, \dots, N\}$ there exists $p = p(i) \in \mathbb{N}$ such that

$$F^{m \times tp}(B(\mathbf{x}^{(i)}, \epsilon_i)) \subset B(\mathbf{x}^{(i)}, \epsilon_i) \quad \forall m \in \mathbb{N}.$$

Contractivity implies that $B(\mathbf{x}^{(i)}, \epsilon_i) \cap \Lambda = \{\mathbf{x}^{(i)}\}$ for each $i \in \{1, 2, \dots, N\}$. Furthermore, $\mathbf{x}^{(i)}$ is periodic of period (not necessarily minimal) $t \times p(i)$. Periods are bounded by $N \times t$, so they depend on $\text{dist}(\Lambda, \Delta)$ through the Lebesgue number of $\mathcal{C}$. $\square$

We would like to prove that, in the case of networks of piecewise affine contractions, $\text{dist}(\Lambda, \Delta) > 0$ is generic. This is true for in the 1-dimensional case [4], but the question remains open for $d \geq 2$.

When on the contrary $\text{dist}(\Lambda, \Delta) = 0$, the attractor may fail to be invariant under the dynamics. In this case, to each point $\mathbf{x} \in \Lambda$ we may associate a “ghost orbits”, which would be a real orbit by a convenient redefinition of the transformation at the discontinuities.

Given an orbit $O(\mathbf{x})$ and $\tau \in \mathbb{N}$, let $\mathcal{P}_{\mathbf{x}}^{\tau} := \{\mathbf{I} \in \mathcal{P}^{\tau} : \exists t \in \mathbb{N} \text{ such that } \mathbf{x}^{(t)} \in \mathbf{I}\}$. This is the restriction of the dynamical partition to the orbit. We will say that the orbits $O(\mathbf{x})$ and $O(\mathbf{y})$ are distinct if $\mathcal{P}_{\mathbf{x}}^{\tau} \cap \mathcal{P}_{\mathbf{y}}^{\tau} = \emptyset$ for some $\tau \in \mathbb{N}$. Note that distinct orbits are necessarily distinguishable at some time. i. e., if $O(\mathbf{x})$ and $O(\mathbf{y})$ are distinct then there exists $\tau \in \mathbb{N}$ such that $\mathbf{x}^{\tau}$ and $\mathbf{y}^{\tau}$ belong to different atoms of the base partition $\mathcal{P}$.

We can easily relate the growth of the dynamical complexity to the number of infinite, distinct orbits. Indeed, for $O(\mathbf{x})$ infinite, we have $\#\mathcal{P}_{\mathbf{x}}^t \geq t + 1$ for all $t \in \mathbb{N}$. Otherwise there would exist $t \in \mathbb{N}$ such that $\mathcal{P}^t = \mathcal{P}^{t+1}$ and the orbit would be ultimately periodic. Hence, if Λ contains K distinct infinite orbits $O(\mathbf{x}^1), O(\mathbf{x}^2), \dots, O(\mathbf{x}^K)$, then there exists a time $\tau \in \mathbb{N}$ such that $C(t) \geq \sum_{i=1}^K \#\mathcal{P}_{\mathbf{x}^i}^t \geq K(t + 1)$ for each $t \geq \tau$. Therefore, assuming the lowest possible complexity for each infinite component of the attractor, a linear behavior of the complexity would imply the finiteness of the number of such components.

The dynamical complexity is directly related to the topological entropy. For a dynamical system (X, F) , for each $\tau \in \mathbb{N}$ and $\epsilon > 0$, a set $E \subset X$ is said to be (τ, ϵ) -separated if for $\mathbf{x}, \mathbf{y} \in E$ there exists $0 \leq t \leq \tau$ such that $d(F^t(\mathbf{x}), F^t(\mathbf{y})) \geq \epsilon$. The topological entropy of the system (X, F) is defined by

$$h_{\text{top}} := \lim_{\epsilon \rightarrow 0} (\limsup_{\tau \rightarrow \infty} \log N(\tau, \epsilon) / \tau),$$

where $N(\tau, \epsilon) = \max\{\#E : E \subset X \text{ is } (\tau, \epsilon)\text{-separated}\}$.

The topological entropy of a piecewise contraction (X, F) satisfies the inequality

$$h_{\text{top}} \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log C(t).$$

Indeed, for $\tau \in \mathbb{N}$, $\epsilon > 0$ and $\mathbf{I} \in \mathcal{P}^{\tau}$ fixed, if $E \subset X$ is (τ, ϵ) -separated, then $E \cap \mathbf{I}$ is necessarily $(1, \epsilon)$ -separated. Therefore $N(\tau, \epsilon) \leq C(\tau) \times N(1, \epsilon)$.

3. MAIN RESULT

Theorem 3.1 (Polynomial upper bound). *Given the interaction and threshold matrices $K, T \in \mathcal{M}_{d \times d}([0, 1])$, and the activation matrix $s \in \mathcal{M}_{d \times d}(\{-1, 0, 1\})$, there exist $a_0 \in [0, 1]$ and $c \geq 1$, such that for each contraction rate $a \in [0, a_0)$, the complexity of corresponding discrete time regulatory network satisfies $C(t) \leq 1 + c(1 + c^d t^d)$.*

In the next section we will consider some specific constraints on the structure of the network with which we are able to improve this upper bound. We will also show some concrete examples where the constant involved in the upper bound is explicitly determined.

3.1. Prerequisites. For each $1 \leq j \leq d$ let us define the system of affine 1-dimensional contractions

$$(3) \quad \mathcal{F}_j := \left\{ x \mapsto ax + (1-a) \left(\sum_{i=1}^d \mathbf{e}_i K_{i,j} \right) : \mathbf{e} \in \{0,1\}^d \right\}.$$

This system of contractions allows us to compute the coordinates of F . Indeed, for each $\mathbf{I} \in \mathcal{P}$ and $1 \leq j \leq d$, there exists an affine contraction $f \in \mathcal{F}_i$ such that $F(\mathbf{x})_j = f(\mathbf{x}_j)$ whenever $\mathbf{x} \in \mathbf{I}$.

The discrete time regulatory network $([0,1]^d, F)$ is said to satisfy *coordinatewise injectivity* if for each $1 \leq j \leq d$ and $f, f' \in \mathcal{F}_j$ we have

$$f([0,1]) \cap f'([0,1]) = \emptyset \quad \text{whenever} \quad f \neq f'.$$

To prove the main result of this paper we first show that for networks satisfying this property, $C(t)$ grows at most polynomially. In particular, for a network satisfying coordinatewise injectivity, the corresponding mapping F is injective. In this case the sets in $\mathcal{Q}^{t+1} := F^t(\mathcal{P}^{t+1})$ are disjoint affine images of the atoms in $\mathcal{P}^t$, and we can use the equation $\#\mathcal{P}^t = \#\mathcal{Q}^t$ to compute the complexity. This equality holds in the more general situation, when for each $t \in \mathbb{N}$, F^t maps different atoms in $\mathcal{P}$ to different sets, not necessarily disjoint.

In the coordinatewise injective case, the degree of the polynomial bound for the complexity may be strictly smaller than the cardinality of the network. In the next section we will see how this degree can be related to the topological structure of the network.

Lemma 3.1. *If $([0,1]^d, F)$ satisfies coordinatewise injectivity, then F is injective.*

Proof. For $\mathbf{x}, \mathbf{y} \in [0,1]^d$ with $\mathbf{x} \neq \mathbf{y}$ we have two possibilities: either they belong to the same atom of the base partition, or they belong to different atoms.

In the first case, since the piecewise constant part of F

$$\mathbf{x} \mapsto (1-a) \left(\sum_{i=1}^d K_{i,j} H(s_{i,j}(\mathbf{x}_i - T_{i,j})) \right)_{j=1}^d$$

is constant on the atoms of $\mathcal{P}$, it follows that $F(\mathbf{x}) - F(\mathbf{y}) = a(\mathbf{x} - \mathbf{y}) \neq 0$.

In the second case there exists $1 \leq i \leq d$, and different affine contractions $f, f' \in \mathcal{F}_i$, such that $F(\mathbf{x})_i = f(\mathbf{x}_i)$ while $F(\mathbf{y})_i = f'(\mathbf{y}_i)$. The coordinatewise injectivity ensures that f and f' have disjoint images, which implies that $F(\mathbf{x})_i \neq F(\mathbf{y})_i$, and hence $F(\mathbf{x}) - F(\mathbf{y}) \neq 0$ as required. $\square$

Proposition 3.1 (Strong contraction implies coordinatewise injectivity). *For each $K \in \mathcal{M}_{d \times d}([0, 1])$ there exists $a_0 > 0$ such that any discrete time regulatory network with interaction matrix K and contraction rate $a \in [0, a_0)$ satisfies coordinatewise injectivity.*

Proof. Fix the threshold and interaction matrices compatible with K . For each $j \in \{1, 2, \dots, d\}$, let $\mathcal{F}_j$ be the system of affine contractions defined in (3). For $j \in \{1, 2, \dots, d\}$ fixed, let

$$\mathcal{I}_j := \left\{ \eta \in [0, 1] : \eta := \sum_{i=1}^d \mathbf{e}_i K_{i,j} \text{ for some } \mathbf{e} \in \{0, 1\}^d \right\}.$$

Let $\delta_j := \min\{|\eta - \eta'| : \eta, \eta' \in \mathcal{I}_j \text{ and } \eta \neq \eta'\}$. Finally, let $\delta := \min_{1 \leq j \leq d} \delta_j$, which is strictly positive. With this define $a_0 := \delta/(1 + \delta)$, so that $a < (1 - a)\delta$ for each $0 \leq a < a_0$.

Suppose that $a < a_0$, fix $i \in \{1, 2, \dots, d\}$, and take $f, f' \in \mathcal{F}_i$ such that $f \neq f'$. Hence there are $\eta, \eta' \in \mathcal{I}_i$ with $\eta \neq \eta'$, such that $f(x) := ax + (1 - a)\eta$ and $f'(x) := ax + (1 - a)\eta'$. Without loss of generality, let us suppose that $\eta > \eta'$, which implies that $\eta \geq \eta' + \delta$. Then we have

$$\max_{x \in [0, 1]} f'(x) = a + (1 - a)\eta' < (1 - a)(\eta' + \delta) < (1 - a)\eta = \min_{x \in [0, 1]} f(x),$$

which holds for $i \in \{1, 2, \dots, d\}$ arbitrary, and the proposition follows. $\square$

For each $1 \leq i \leq d$ and $t \in \mathbb{N}$, let $\mathcal{Q}_i^t$ be the collection of intervals

$$\mathcal{Q}_i^t := \{J \subset [0, 1] : J = \Pi_i \mathbf{J} \text{ for some } \mathbf{J} \in \mathcal{Q}^t\},$$

Let us recall that $\mathcal{Q}^{t+1}$ is the image, at time t , of the dynamical partition $\mathcal{P}^{t+1}$. Here and below Π_i will denote the projection on the i -th coordinate.

Lemma 3.2. *If $([0, 1]^d, F)$ satisfies coordinatewise injectivity, then for each $1 \leq i \leq d$ and $t \in \mathbb{N}$, the intervals in $\mathcal{Q}_i^t$ are pairwise disjoint.*

Proof. For each $1 \leq i \leq d$ $\mathcal{Q}_i^1 = \mathcal{P}_i$ is a partition, and so the intervals in $\mathcal{Q}_i^1$ are pairwise disjoint. Suppose that for $t = \tau$ the intervals in $\mathcal{Q}_i^\tau$ are pairwise disjoint. Let $\mathcal{F}_i$ be the system of affine contractions as defined in equation (3). Then

$$\mathcal{Q}_i^{\tau+1} \subset \mathcal{F}_i(\mathcal{Q}_i^\tau) \vee \mathcal{P}_i := \{f(J) \cap I : f \in \mathcal{F}_i, J \in \mathcal{Q}_i^\tau, \text{ and } I \in \mathcal{P}_i\}.$$

Since F satisfies coordinatewise injectivity, and since the intervals in $\mathcal{Q}_i^\tau$ are mutually disjoint, it follows that $\mathcal{F}_i(\mathcal{Q}_i^\tau) \vee \mathcal{P}_i$ is a collection of pairwise disjoint intervals, implying that the intervals in $\mathcal{Q}_i^{\tau+1}$ are pairwise disjoint as well. $\square$

Given a set of coordinates $U \subset \{1, 2, \dots, d\}$, and $t \in \mathbb{N}$ fixed, a (U, t) -specification is a $\#U$ -dimensional interval $\mathbf{S} \in \prod_{i \in U} \mathcal{Q}_i^t$. The *bouquet* corresponding to the specification $\mathbf{S}$ is the collection of d -dimensional intervals

$$(4) \quad C(\mathbf{S}) := \{\mathbf{J} \in \mathcal{Q}^t : \Pi_i \mathbf{J} = \Pi_i \mathbf{S} \ \forall i \in U\}.$$

We will say that $\mathbf{J} \in \mathcal{Q}^t$ satisfies specification $\mathbf{S}$ if $\mathbf{J} \in C(\mathbf{S})$, and we will call $N(\mathbf{S}) := \#C(\mathbf{S})$ the *degeneracy* of $\mathbf{S}$.

Corollary 3.1. *If $([0, 1]^d, F)$ satisfies coordinatewise injectivity, for each $1 \leq i, j \leq d$ and $t \in \mathbb{N}$, there exists an interval $J_{i,j}^t \in \mathcal{Q}_i^t$ such that $\{\mathbf{J} \in \mathcal{Q}^t : T_{i,j} \in \Pi_i F(\mathbf{J})\} \subset C(J_{i,j}^t)$.*

Proof. For each $\mathbf{J} \in \mathcal{Q}^t$ we have $\Pi_i F(\mathbf{J}) \in \mathcal{F}_i(\mathcal{Q}_i^t) := \{f(J) : f \in \mathcal{F}_i \text{ and } J \in \mathcal{Q}_i^t\}$. The previous lemma ensures that $\mathcal{F}_i(\mathcal{Q}_i^t)$ consists of partwise disjoint intervals. Thus, for each $(i, j) \in \mathcal{A}$, there cannot be more than one interval $J \in \mathcal{Q}^t$, and one affine contraction $f \in \mathcal{F}_i$, such that $T_{i,j} \in f(J)$. $\square$

We denote $J_{i,j}^t$ the unique interval in $\mathcal{Q}_i^t$ such that if $T_{i,j} \in \Pi_i F(\mathbf{J})$ for $\mathbf{J} \in \mathcal{Q}^t$, then $\Pi_i \mathbf{J} = J_{i,j}^t$.

We say that the d -dimensional interval $\mathbf{J} \in \mathcal{Q}^t$ is a *predecessor* of $\mathbf{J}' \in \mathcal{Q}^{t+1}$ if $\mathbf{J}' = F(\mathbf{J}) \cap \mathbf{I}$ for some $\mathbf{I} \in \mathcal{P}$. Reciprocally we say that $\mathbf{J}'$ is a *successor* of $\mathbf{J}$. In our case, since F is injective, every $\mathbf{J}' \in \mathcal{Q}^{t+1}$ has a unique predecessor $\mathbf{J} \in \mathcal{Q}^t$, which we denote by $P(\mathbf{J}')$.

Lemma 3.3. *Suppose that $([0, 1]^d, F)$ is coordinatewise injective. Fix $U \subset \{1, 2, \dots, d\}$ and $t \in \mathbb{N}$. For each $(U, t+1)$ -specification $\mathbf{S}$, there is a unique (U, t) -specification $P(\mathbf{S})$, such that $\mathbf{J} \in C(\mathbf{S})$ implies $P(\mathbf{J}) \in C(P(\mathbf{S}))$.*

Proof. Fix $i \in U$ and $\mathbf{J} \in \mathcal{Q}^{t+1}$ satisfying the specification $\mathbf{S}$. Let $\mathbf{J}' = P(\mathbf{J})$ and $\mathbf{I} \in \mathcal{P}$ such that $\mathbf{J} = F\mathbf{J}' \cap \mathbf{I}$. Then

$$(5) \quad \Pi_i \mathbf{S} = \Pi_i (F(\mathbf{J}') \cap \mathbf{I}) = f(\Pi_i \mathbf{J}') \cap \Pi_i \mathbf{I}$$

for some $f \in \mathcal{F}_i$. Since F is coordinatewise injective, then the affine contraction $f \in \mathcal{F}_i$ for which (5) holds is uniquely determined from $\mathbf{J}' \in \mathcal{Q}^t$. On the other hand, Lemma 3.2 establishes that for each $1 \leq i \leq d$ and $t \in \mathbb{N}$, the collection $\mathcal{Q}_i^t$ consists of partwise disjoint intervals. Hence, for $f \in \mathcal{F}_i$ fixed, there is only one interval $J' \in \mathcal{Q}_i^t$ such that $\Pi_i \mathbf{J}' = J'$ satisfies Equation (5). Therefore $\Pi_i \mathbf{J}'$ depends only on $\Pi_i \mathbf{S}$, and the lemma is proved. $\square$

Lemma 3.4. *Let $([0, 1]^d, F)$ be coordinatewise injective, and let $U \subset \{1, 2, \dots, d\}$. For each $t \in \mathbb{N}$, and every $(U, t+1)$ -specification $\mathbf{S}$, we have*

$$N(\mathbf{S}) \leq N(P(\mathbf{S})) + c \times \sum_{i \notin U} \sum_{j=1}^d N(P(\mathbf{S}) \times J_{i,j}^t),$$

with J_j^t as defined by above, and $c := \#\mathcal{P} - 1$.

Proof. Lemma 3.3 implies that

$$C(\mathbf{S}) = \{F(\mathbf{J}') \cap \mathbf{I} \in C(\mathbf{S}) : \mathbf{I} \in \mathcal{P} \text{ and } \mathbf{J}' \in C(P(\mathbf{S}))\}.$$

If $\mathbf{J}' \in C(P(\mathbf{S}))$ has more than one successor satisfying $\mathbf{S}$, then necessarily $T_{i,j} \in \Pi_i F(\mathbf{J}')$ for some $i \notin U$ and $1 \leq j \leq d$. Now, each $\mathbf{J}' \in C(P(\mathbf{S}))$ has no more than $\#\mathcal{P}$ successors. Therefore

$$N(\mathbf{S}) \leq N(P(\mathbf{S})) + c \times \#\{\mathbf{J}' \in C(P(\mathbf{S})) : T_{i,j} \in \Pi_i F(\mathbf{J}') \text{ for some } i \notin U \text{ and } 1 \leq j \leq d\}.$$

From Lemma 3.1, if $\mathbf{J}' \in C(P(\mathbf{S}))$ has more than one successor that satisfies $\mathbf{S}$, then $\mathbf{J}'$ satisfies $P(\mathbf{S}) \times J_{i,j}^t$ for some $i \notin U$ and $1 \leq j \leq d$. Thus,

$$\{\mathbf{J}' \in C(P(\mathbf{S})) : T_{i,j} \in \Pi_i F(\mathbf{J}') \text{ for some } i \notin U \text{ and } 1 \leq j \leq d\} \subset \bigcup_{i \notin U} \bigcup_{j=1}^d C(P(\mathbf{S}) \times J_{i,j}^t),$$

and from this we readily obtain the required inequality. $\square$

3.2. Proof of Theorem 3.1. Fix $a \in [0, a_0)$, with $a_0 > 0$ depending on $K, T \in \mathcal{M}_{d \times d}([0, 1])$ and $s \in \mathcal{M}_{d \times d}(\{-1, 0, 1\})$ as in the proof of Proposition 3.1. This proposition ensures that the mapping F satisfies coordinatewise injectivity. Lemma 3.1 ensures that F is injective, and so we can replace $\mathcal{P}^t$ by $\mathcal{Q}^t$ in the computation of the complexity. In order to keep control of the growth of $\#\mathcal{Q}^t$, we use the recurrence $\mathcal{Q}^{t+1} = F(\mathcal{Q}^t) \vee \mathcal{P}$.

If $C(t+1) > C(t)$, then the additional contribution in $C(t+1)$ is due to those d -dimensional intervals in $\mathcal{Q}^t$ which posses more than one successor.

Since $\mathbf{J} \in \mathcal{Q}^t$ cannot have more than $\#\mathcal{P}$ successors in $\mathcal{Q}^{t+1}$, one for each possible intersection $F(\mathbf{J}) \cap \mathbf{I}$ with $\mathbf{I} \in \mathcal{P}$, it follows that the complexity satisfies the recursion

$$(6) \quad C(t+1) \leq C(t) + c_1 \times \#\{\mathbf{J} \in \mathcal{Q}^t : \mathbf{J} \text{ has more than two successors}\},$$

where $c := \#\mathcal{P} - 1$.

Now, for $\mathbf{J} \in \mathcal{Q}^t$ to have more than one successor it is required that $F(\mathbf{J})$ intersect at least two different atoms in $\mathcal{P}$. But if this is the case, then necessarily $F(\mathbf{J})$ contains a discontinuity set for F . Hence, if $\mathbf{J} \in \mathcal{Q}^t$ has more than one successor, then there exists $1 \leq i, j \leq d$ such that $T_{i,j} \in \Pi_i F(\mathbf{J})$. Then, according to (6) we have

$$(7) \quad C(t+1) \leq C(t) + c \times \#\{\mathbf{J} \in \mathcal{Q}^t : T_{i,j} \in \Pi_i F(\mathbf{J}) \text{ for some } 1 \leq i, j \leq d\}.$$

Using lemma 3.1 and (7) we conclude that

$$C(t+1) \leq C(t) + c \times \#\{\mathbf{J} \in \mathcal{Q}^t : \Pi_i \mathbf{J} = J_{i,j}^t \text{ for some } 1 \leq i, j \leq d\},$$

and so we can rewrite (7) as

$$(8) \quad C(t+1) \leq C(t) + c \times \sum_{i=1}^d \sum_{j=1}^d \#\{\mathbf{J} \in \mathcal{Q}^t : \Pi_i \mathbf{J} = J_{i,j}^t\}.$$

Since the cardinalities $\#\{\mathbf{J} \in \mathcal{Q}^t : \Pi_j \mathbf{J} = J_{i,j}^t\}$ in (8) are the same as the degeneracies $N(J_{i,j}^t)$, we can rewrite (8) as

$$(9) \quad C(t+1) \leq C(t) + c \times \sum_{i=1}^d \sum_{j=1}^d N(J_{i,j}^t),$$

where $c := \#\mathcal{P} - 1$.

In what follows we will establish polynomial bounds for the growth of the degeneracies $N(J)$, with $J \in \mathcal{Q}_i^t$ arbitrary. For this aim we use a recurrence relation between degeneracies of successive specifications. According to Lemma 3.3, for $U \subset \{1, 2, \dots, d\}$ and $t \in \mathbb{N}$ fixed, to each $(U, t+1)$ -specification $\mathbf{S}$, there corresponds a unique (U, t) -specification $P(\mathbf{S})$, such that if $\mathbf{J} \in \mathcal{Q}^{t+1}$ satisfies $\mathbf{S}$ then its predecessor $P(\mathbf{J})$ satisfies $P(\mathbf{S})$. Then Lemma 3.4 establishes the inequalities

$$(10) \quad N(\mathbf{S}) \leq N(P(\mathbf{S})) + c \times \sum_{i \notin U} \sum_{j=1}^d N(P(\mathbf{S}) \times J_{i,j}^t),$$

satisfied for each $(U, t+1)$ -specification $\mathbf{S}$, with $U \subset \{1, 2, \dots, d\}$. We do not solve this chain of inequalities, but instead we solve the chain corresponding to the maximal degeneracies. Indeed, for each $U \subset \{1, 2, \dots, d\}$ and $t \in \mathbb{N}$, let

$$n(U, t) := \max\{N(\mathbf{S}) : \mathbf{S} \text{ is a } (U, t)\text{-specification}\}.$$

Taking the maximum at both sides of the inequality (10), we obtain the chain of recursions

$$(11) \quad n(U, t+1) \leq n(U, t) + c \times \sum_{i \notin U} (\#\mathcal{P}_i - 1) \times n(U \cup \{i\}, t)$$

taking into account that $\#\{1 \leq j \leq d : T_{i,j} > 0\} = \#\mathcal{P}_i - 1$. We solved this chain of recursions as follows.

Let us first take $U := \{1, 2, \dots, d\} \setminus \{i\}$, for $1 \leq i \leq d$ arbitrary. Taking into account that $n(\{1, 2, \dots, d\}, t) = 1$ for each $t \in \mathbb{N}$, from (11) we obtain the linear recursion

$$n(U, t+1) \leq n(U, t) + c \times (\#\mathcal{P}_i - 1).$$

where $c := \#\mathcal{P} - 1$. This recursion has solution $n(U, t) \leq n(U, 1) + c \times (\#\mathcal{P}_i - 1) \times (t - 1)$. Since $n(U, 1) \leq \#\mathcal{P}_i \leq c \times (\#\mathcal{P}_i - 1)$, we have

$$n(U, t) \leq c \times (\#\mathcal{P}_i - 1) \times t, \quad \forall t \in \mathbb{N}.$$

Suppose that for each $U \subset \{1, 2, \dots, d\}$ with $\#\{1 \leq i \leq d : i \notin U\} = k$. Then,

$$(12) \quad n(U, t) \leq c^k \times t^k \times \prod_{i \notin U} (\#\mathcal{P}_i - 1) \quad \forall t \in \mathbb{N}.$$

Let $U' \subset \{1, 2, \dots, d\}$, with $\#\{1 \leq i \leq d : i \notin U'\} = k + 1$. Substituting this into (11), and taking into account the induction hypothesis, we obtain

$$\begin{aligned} n(U', t + 1) &\leq n(U', t) + c \times \sum_{i \notin U'} (\#\mathcal{P}_i - 1) \times n(U' \cup \{i\}, t) \\ &\leq n(U', t) + (k + 1) c^{k+1} \times t^k \times \prod_{j \notin U'} (\#\mathcal{P}_j - 1), \end{aligned}$$

with solution $n(U', \tau) \leq n(U', 1) + (k + 1) c^{k+1} \times \left(\sum_{t=1}^{\tau-1} t^k \right) \times \prod_{j \notin U'} (\#\mathcal{P}_j - 1)$, for each $\tau \in \mathbb{N}$.

Now, taking into account that $\sum_{t=1}^{\tau-1} t^k \leq \tau^{k+1}/(k + 1)$, we get

$$n(U', \tau) \leq c^{k+1} \times \tau^{k+1} \times \prod_{j \notin U'} (\#\mathcal{P}_j - 1),$$

which proves inequality (12) for each $U \subset \{1, 2, \dots, d\}$.

Since $N(J) \leq n(\{j\}, t)$ for each $t \in \mathbb{N}$ and every $J \in \mathcal{Q}_j^t$, by replacing (12) into (9), we obtain an autonomous recursion satisfied by the complexity:

$$C(t + 1) \leq C(t) + c \times dc^{d-1} \times t^{d-1} \times \prod_{i=1}^d (\#\mathcal{P}_i - 1).$$

We solve this recursion taking into account that $C(1) = \#\mathcal{P} = c + 1$ and that $\prod_{i=1}^d (\#\mathcal{P}_i - 1) \leq c$, and finally obtain the upper bound

$$\begin{aligned} C(\tau) &\leq 1 + c \left(1 + dc^d \left(\sum_{t=1}^{\tau-1} t^{d-1} \right) \right) \\ &\leq 1 + c \left(1 + c^d \tau^d \right), \end{aligned}$$

for each $\tau \in \mathbb{N}$. □

3.3. Generalizations. Theorem 3.1 admits two quite direct generalizations as follows.

Rectangular piecewise contractions. Lemmas from 3.1 to 3.4 can be extended to piecewise contractions of the following type: give the base partition $\mathcal{P} := \prod_{i=1}^d \mathcal{P}_i$, let $F : [0, 1]^d \rightarrow [0, 1]^d$ be such that for each $1 \leq i \leq d$ and $\mathbf{I} \in \mathcal{P}$, there exists a contractive map $f_{i,\mathbf{I}} : [0, 1] \rightarrow [0, 1]$ such that

$$F(\mathbf{x})_i = f_{i,\mathbf{I}}(\mathbf{x}_i), \quad \forall \mathbf{x} \in \mathbf{I}.$$

We say that F so defined is coordinatewise injective when for each $1 \leq i \leq d$, if $f_{i,\mathbf{I}} \neq f_{i,\mathbf{J}}$ then $f_{i,\mathbf{I}}([0, 1]) \cap f_{i,\mathbf{J}}([0, 1]) = \emptyset$. Theorem 3.1 holds for systems $([0, 1]^d, F)$, where F is a piecewise contractions of the type just described, satisfying the coordinatewise injectivity.

We say that a piecewise contraction $F : [0, 1]^d \rightarrow [0, 1]^d$ of the type describe above is a *rectangular piecewise contraction* compatible with $\mathcal{P}$. We use the adjective rectangular since such a transformation, when restricted to an atom of $\mathfrak{P}$, maps the products of intervals into products of intervals.

The bound $C(t) \leq 1 + c(1 + c^d t^d)$ holds for any d -dimensional regulatory network, as far as the piecewise injectivity is ensured. Hence, lemma 3.1 implies that given a interaction matrix M , all the regulatory networks with interaction matrix M and sufficiently small contraction rate, have complexity bounded by the same polynomial $t \mapsto 1 + c(1 + c^d t^d)$.

Compatible sequences. Instead of a single rectangular piecewise contraction $F : [0, 1]^d \rightarrow [0, 1]^d$ compatible with a given base partition $\mathcal{P}$, we may consider a sequence $SS := \{F_t : [0, 1]^d \rightarrow [0, 1]^d\}_{t=1}^\infty$ of rectangular piecewise contractions compatible with $\mathcal{P}$. Then, we may consider the non-autonomous system generated by SS where orbits are given by

$$\mathcal{O}_{SS}(\mathbf{x}) := \{\mathbf{x}^t := F_t(\mathbf{x}^{t-1}), \mathbf{x}^0 = \mathbf{x}\}.$$

The *dynamical complexity of the sequence SS* is the integer function $C_{SS} : \mathbb{N} \rightarrow \mathbb{N}$ such that $C_{SS}(t) := \#\mathcal{P}_{SS}^t$, where $\mathcal{P}_{SS}^1 := \mathcal{P}$ and $\mathcal{P}_{SS}^{t+1} := \{F_{t+1}^{-1}(\mathbf{I}) \cap \mathbf{J} : \mathbf{I} \in \mathcal{P}_{SS}^t \text{ and } \mathbf{J} \in \mathcal{P}\}$, for $t \geq 1$. If each one of the maps in SS satisfies the coordinatewise injectivity, then lemmas 3.1 to 3.4 hold. In this case we use $\mathcal{Q}_{SS}^t := F_t \circ \dots \circ F_2 \circ F_1(\mathcal{P}_{SS}^t)$ in place of $\mathcal{Q}^t$.

Mutatis mutandi one can prove the analogous of Theorem 3.1.

Theorem 3.2. *Let $SS := \{F_t : [0, 1]^d \rightarrow [0, 1]^d\}_{t=1}^\infty$ be sequence of rectangular piecewise contractions compatible with a partition $\mathcal{P} := \prod_{i=1}^d \mathcal{P}_i$. If all the contractions in SS satisfy the coordinatewise injectivity, then $C_{SS}(t) \leq 1 + c(1 + c^d t^d)$ with $c := \#\mathcal{P} - 1$.*

In what follows a sequence of rectangular piecewise contractions compatible with a given partition $\mathcal{P}$, all of them satisfying the coordinatewise injectivity, will be refereed as a $\mathcal{P}$ -compatible sequence.

Remark 3.1. *The complexity of a compatible sequence of piecewise contractions with base partition $\mathcal{P}$ is bounded by the polynomial $t \mapsto 1 + c(1 + c^d t^d)$. Note that the constant c depends only on the cardinality of the base partition, indeed, $c := \#\mathcal{P} - 1$.*

4. THE UNDERLYING NETWORK

A *network* (directed graph) consist on a finite set V of *vertices*, and a set of ordered couples $\mathcal{A} \subset V \times V$, the *arrows*. We consider the interacting units to form a network with vertex set $V := \{1, 2, \dots, d\}$ and arrow set $\mathcal{A} := \{(i, j) \in V \times V : K_{i,j} > 0\}$. We call this *the underlying network*. As

we mentioned above, the main motivation of this study is to understand the relationship between the structure of the underlying network and the asymptotic dynamics of the system.

Let $T, K : \{1, 2, \dots, d\} \times \{1, 2, \dots, d\} \rightarrow [0, 1]$ and $s : \{1, 2, \dots, d\} \times \{1, 2, \dots, d\} \rightarrow \{-1, 0, 1\}$ be compatible matrices. Assume that for each $1 \leq j \leq d$, $\sum_{i=1}^d K_{i,j} < 1$ and fix a contraction rate $a \in (0, 1)$. To the sequence of vectors $\mathcal{D} := \{D_t \in [0, 1]^d\}_{t=1}^\infty$ satisfying $0 \leq D_{t,j} + \sum_{i \in V} K_{i,j} \leq 1$, we associate the sequence $\mathcal{D}_{K,T,s,a} := \{F_t : [0, 1]^d \rightarrow [0, 1]^d\}_{t=1}^\infty$ of piecewise affine contractions such that

$$F_t(\mathbf{x})_j := a\mathbf{x}_j + (1-a) \left(\sum_{i \in V} K_{i,j} H(s_{i,j}(T_{i,j} - \mathbf{x}_i)) + D_{t,j} \right),$$

for each $t \in \mathbb{N}$ and $1 \leq i \leq d$. Notice that threshold matrix T completely determines the partition $\mathcal{P} := \prod_{i \in V} \mathcal{P}_i$ whose atoms are the domains of continuity of the maps $F_t \in \mathcal{D}_{K,T,s,a}$. Taking $a \in (0, 1)$ sufficiently small so that each transformation $F_t \in \mathcal{D}_{K,T,s,a}$ is coordinatewise injective, we obtain a $\mathcal{P}$ -compatible sequence. In this way we can define the complexity

$$(13) \quad C_{T,K,s,a}(t) := \max \left\{ C_{\mathcal{D}_{T,K,s,a}}(t) : \mathcal{D} \text{ such that } 0 \leq D_{t,j} + \sum_{i \in V} K_{i,j} \leq 1 \forall t \in \mathbb{N} \ j \in V \right\}$$

for the parameters K, T, s, a .

4.1. Skew product. The first result in this direction concerns networks where the dynamics on a subnetwork is forced by the dynamics on the complementary subnetwork. Let (V, A) be a network such that the vertex set V may be decomposed into two disjoint sets $V = V_{\text{base}} \cup V_{\text{bundle}}$ such that $A \cap (V_{\text{bundle}} \times V_{\text{base}}) = \emptyset$. We say in this case that (V, A) admits *base–bundle decomposition*.

When the underlying network (V, A) of the system $([0, 1]^{\#V}, F)$ admits a base–bundle decomposition $V = V_b \cup V_d$, the associate interaction matrix $K \in \mathcal{M}_{\#V}([0, 1])$ has the triangular block form

$$K = \begin{pmatrix} K_{b,b} & K_{b,d} \\ \mathbf{0} & K_{d,d} \end{pmatrix},$$

where $K_{b,b} := K|_{V_b \times V_b}$, $K_{b,d} := K|_{V_b \times V_d}$ and $K_{d,d} := K|_{V_d \times V_d}$. In this case the projection $F(\mathbf{x})_b := \{\Pi_j F(\mathbf{x}) : j \in V_b\}$ depends only on $\mathbf{x}_b := \{\Pi_j \mathbf{x} : j \in V_b\}$, thus the projection of $([0, 1]^{\#V}, F)$ to coordinates in V_{base} is well defined.

If the underlying network (V, A) of the system $([0, 1]^{\#V}, F)$ admits a base–bundle decomposition $V = V_b \cup V_d$, then the dynamics has the structure of a *skew product*. Indeed, let

$$F_b : [0, 1]^{\#V_b} \rightarrow [0, 1]^{\#V_b} \text{ such that } F_b(\mathbf{y}) := F(\mathbf{y} \oplus \mathbf{z})$$

with $\mathbf{z} \in [0, 1]^{\#V_d}$ arbitrarily chosen, and for each $\mathbf{y} \in [0, 1]^{\#V_b}$ let

$$F_{\mathbf{y}} : [0, 1]^{\#V_b} \rightarrow [0, 1]^{\#V_b} \text{ such that } F_{\mathbf{y}}(\mathbf{z}) := F(\mathbf{y} \oplus \mathbf{z}).$$

Then, for $\mathbf{x} \equiv \mathbf{x}_b \oplus \mathbf{x}_d \in [0, 1]^d$, we have $F(\mathbf{x}) = F_b(\mathbf{x}_b) \oplus F_{\mathbf{x}_b}(\mathbf{x}_d)$.

Theorem 4.1. *For a regulatory network with parameters T, K, s, a , suppose that the associated underlying network (V, A) admits a base-bundle decomposition $V = V_b \cup V_d$. If $([0, 1]^{\#V}, F)$ satisfies coordinatewise injectivity, then*

$$C(t) \leq C_b(t) \times C_{(K_d, T_d, s_d, a)}(t),$$

where $t \mapsto C_b(t)$ is the dynamical complexity of the regulatory network $([0, 1]^{\#V_b}, F_b)$, and K_d, T_d and s_d denote respectively the restrictions of K, T and s to $V_d \times V_d$.

Proof. Let $\mathcal{P}_b := \prod_{i \in V_b} \mathcal{P}_i$ and $\mathcal{P}_d := \prod_{i \in V_d} \mathcal{P}_i$ be respectively the projection of the base partition to the coordinates in V_b and V_d . Because of the skew product structure of the map, for each $t \in \mathbb{N}$ and $\mathbf{I} \in \mathcal{P}^t$ we have intervals $\mathbf{I}_b \in \mathcal{P}_b^t$ and $\mathbf{I}_d \in [0, 1]^{\#V_d}$ such that $\mathbf{I} = \mathbf{I}_b \times \mathbf{I}_d$. Since for each $\mathbf{x}_b \in [0, 1]^{\#V_b}$, the corresponding bundle transformation $F_{\mathbf{x}_b} : [0, 1]^{\#V} \rightarrow [0, 1]^{\#V}$ depends only on the atom $\mathbf{J}_b \in \mathcal{P}_b$ containing $\mathbf{x}_b$, then for each $\mathbf{I}_b \in \mathcal{P}_b^t$ there exists a finite sequence of piecewise transformations $F_1, F_2, \dots, F_t : [0, 1]^{\#V_d} \rightarrow [0, 1]^{\#V_d}$ such

$$F^t(\mathbf{I}_b \times \mathbf{I}_d) = F_t \circ F_{t-1} \circ \dots \circ F_1(\mathbf{I}_d),$$

for each multidimensional interval $\mathbf{I}_d \subset [0, 1]^{\#V_d}$. Now, since F is coordinatewise injective, then the finite sequence $F_1, F_2, \dots, F_t$ can be seen as the prefix of a $\mathcal{P}_d$ -compatible sequence in the collection $\mathcal{D}^{K_b, T_b, s_d, a}$ for an appropriated sequence of vectors $\mathcal{D}$. From this it follows that

$$N(\mathbf{I}_b) := \#\{\mathbf{I} \in \mathcal{P}^t : \mathbf{I} = \mathbf{I}_b \times \mathbf{I}_d, \text{ for some } \mathbf{I}_d \in [0, 1]^{\#V_d}\} \leq C_{K_d, T_d, s_d, a}(t),$$

for each $\mathbf{I}_b \in \mathcal{P}_b^t$. From this it follows that

$$C(t) \equiv \#\mathcal{P}^t \leq \#\mathcal{P}_b^t \times C_{K_d, T_d, s_d, a}(t) = C_b(t) \times C_{K_d, T_d, s_d, a}(t),$$

and the theorem is proved. $\square$

4.2. Redundant and essential vertices. Theorem 3.1 establishes that the complexity of a discrete time regulatory network, when the contraction rate is sufficiently strong, grows at most polynomially. The degree of this polynomial bound is never larger than the cardinality of the vertex set. In this section we will see how this degree can be related to the structure of the underlying network, via the notions of redundant and essential sets of vertices.

For $i, j \in V = \{1, 2, \dots, d\}$ fixed, for each $t \in \mathbb{N}$, and for every $J \in \mathcal{Q}_i^t$, let

$$(14) \quad C_j(J) = \Pi_j C(J) := \{\Pi_j \mathbf{J} : \mathbf{J} \in \mathcal{Q}^t \text{ and } \Pi_i \mathbf{J} = J\}.$$

In other words, if the i -th projection of $\mathbf{J} \in \mathcal{Q}^t$ is given, i. e., if $\mathbf{J}$ satisfies a given $(\{i\}, t)$ -specification, and it is equal to $J \in \mathcal{Q}_i^t$, then its j -th projection belongs to the set $C_j(J) \subset \mathcal{Q}_j^t$. We say that i drives j if the cardinality of $C_j(J)$ is uniformly bounded as J ranges over $\cup_{t=1}^{\infty} \mathcal{Q}_i^t$, i. e., if there exists a constant $m_{i,j} \in \mathbb{N}$ such that $\#C_j(J) \leq m_{i,j}$ for all $J \in \cup_{t=1}^{\infty} \mathcal{Q}_i^t$.

A set of vertices $U \subset V$ is *redundant* if for each $i \in U$ there exists $j \in V \setminus U$ such that i drives j in the sense stated above. Below we will see why the driving vertex is the redundant one.

On the other hand, a set $W \subset V$ is *essential* if there exists a fixed multiplicity $M \in \mathbb{N}$ such that for each $t \in \mathbb{N}$ and $\mathbf{S} \in \prod_{i \in W} \mathcal{Q}_i^t$, $N(\mathbf{S}) \leq M$, i. e. $W \subset V$ is essential if the degeneracy of the (W, t) -specifications is uniformly bounded as t varies. Of course $W = V$ is essential, but the interesting case arises when there is a essential proper subset of the vertex set, whose complement is redundant.

Theorem 4.2 (Bounding the degree). *Let $([0, 1]^d, F)$ be a discrete time regulatory network satisfying coordinatewise injectivity, and let $(V, \mathcal{A})$ be the underlying network. Suppose that $U \subset V$ is redundant, whereas $W := V \setminus U$ is essential. Then, there are constants constant $c_1, c_2 > 0$ such that for each $t \in \mathbb{N}$ $C(t) \leq 1 + c_1(1 + c_2 t^q)$, with $q := \#W$.*

The proof of this theorem follows the same scheme, with the suitable modifications, as the proof of Theorem 3.1.

Proof. Since we are assuming coordinatewise injectivity, Lemma 3.1 ensures that F is injective, so that $C(t) = \#\mathcal{Q}^t$. As in the proof of Theorem 3.1, we use the recursion in (6), which can be written as

$$C(t+1) \leq C(t) + c_1 \times \#\{\mathbf{J} \in \mathcal{Q}^t : \Pi_i \mathbf{J} = J_{i,j}^t \text{ for some } (i, j) \in \mathcal{A}\},$$

with $J_{i,j}^t$ as defined by Corollary 3.1.

From here on we use the notion of redundant and essential vertices. Fix $U \subset V$ redundant, such that $W := V \setminus U$ is essential. Now, since each essential vertex can be driven by several redundant ones, let us fix a partition $\mathcal{C} := \{R_j \subset U : j \in W\}$ of U , where to each atom of the partition there corresponds an essential vertex, such that i drives j for each $i \in R_j$. For the rest of the essential vertices let $R_j = \emptyset$. Such partition is not unique since different essential vertices may be driven by the same redundant one. For $\mathcal{C}$ fixed, for each $t \in \mathbb{N}$ and $j \in W$ define the collection of intervals

$$(15) \quad \mathcal{R}_j^t := \bigcup_{i \in R_j} \bigcup_{(i,k) \in \mathcal{A}} C_i(J_{i,k}^t),$$

with $C_i(J_{i,k}^t)$ as defined by (14).

The vertices in R_j are redundant and $\#\{k \in V : (i, k) \in \mathcal{A}\} \leq \mathcal{P}_i - 1$. Therefore

$$(16) \quad \#\mathcal{R}_j^t \leq m_j := (\#\mathcal{P}_j - 1) + \sum_{i \in R_j} m_{i,j} \times (\#\mathcal{P}_i - 1)$$

for each $t \in \mathbb{N}$. Here $m_{i,j}$ is the multiplicity bounding the cardinality of $C_i(J_{i,j}^t)$. For a fixed partition $\mathcal{C}$ and taking into account inequality (6), we obtain

$$(17) \quad C(t+1) \leq C(t) + c_1 \times \sum_{j \in W} \sum_{J \in \mathcal{R}_j^t} N(J),$$

Let us recall that $N(J) := \#C(J) = \#\{\mathbf{J} \in \mathcal{Q}^t : \Pi_i \mathbf{J} = J\}$.

As in the proof of Theorem 3.1, we can find polynomial bounds for the growth of the degeneracies $N(J)$. Taking into account the fact that redundant vertices drive essential ones, we may rewrite inequality (10) as

$$N(\mathbf{S}) \leq N(P(\mathbf{S})) + c_1 \times \sum_{j \in W \setminus W'} \sum_{J \in \mathcal{R}_j^t} N(P(\mathbf{S}) \times J),$$

satisfied for each $(W', t+1)$ -specification $\mathbf{S}$, with $W' \subset W$. Once again we solve the chain of inequalities satisfied by the maximal degeneracies,

$$(18) \quad n(W', t+1) \leq n(W', t) + c_1 \times \sum_{j \in W \setminus W'} m_j n(W' \cup \{j\}, t).$$

Using exactly the same scheme as in the proof of Theorem 3.1 we obtain

$$(19) \quad n(W', t) \leq M c_1^k t^k \prod_{j \in W \setminus W'} n_j, \quad \forall t \in \mathbb{N},$$

with $n_j := \max(m_j, \#\mathcal{P}_j)$.

Since $N(J) \leq n(\{j\}, t)$ for each $t \in \mathbb{N}$ and every $J \in \mathcal{Q}_j^t$, substituting inequality (19) into (17), we obtain

$$\begin{aligned} C(t+1) &\leq C(t) + c_1 \sum_{j \in W} \sum_{J \in \mathcal{R}_j^t} M c_1^{q-1} t^{q-1} \prod_{i \in W \setminus \{j\}} n_i \\ &\leq C(t) + M c_1^q t^{q-1} \sum_{j \in W} m_j \prod_{i \in W \setminus \{j\}} n_j, \end{aligned}$$

with $q = \#W$. We solve this recursion by taking into account that $n_j \geq m_j$ and $C(1) = c_1 + 1$, to finally obtain the upper bound

$$\begin{aligned} C(\tau) &\leq 1 + c_1 \left(1 + q M c_1^q \prod_{j \in W} n_j \left(\sum_{t=1}^{\tau-1} t^{q-1} \right) \right) \\ &\leq 1 + c_1 \left(1 + M c_1^q \tau^q \prod_{j \in W} n_j \right), \end{aligned}$$

for each $\tau \in \mathbb{N}$. The theorem follows by making $c_2 = M c_1^q \prod_{j \in W} n_j$. $\square$

4.3. Head-independent sets. Now we will exhibit concrete situations where the underlying network $(V, \mathcal{A})$ admits a non-trivial redundant vertex set with essential complement. In this way we obtain concrete examples of discrete time regulatory networks whose dynamical complexity is bounded by a polynomial of degree strictly smaller than $\#V$.

Let us first introduce a characterization of a class of vertices which are essential whenever coordinatewise injectivity holds.

Fix a network $(V, \mathcal{A})$. A set of vertices $U \subset V := \{1, 2, \dots, d\}$ is *head-independent* if

- 1) $\{j \in V : (i, j) \in \mathcal{A}\} \cap U = \emptyset$ for all $i \in U$, and
- 2) $\{j \in V : (i, j) \in \mathcal{A}\} \cap \{j \in V : (i', j) \in \mathcal{A}\} = \emptyset$ for all $i, i' \in U$ with $i \neq i'$.

In other words, $U \subset V$ is head-independent if 1) each arrow with tail in U has head outside U , and 2) different arrows with tail in U have necessarily different heads. We will show that for discrete time regulatory network satisfying coordinatewise injectivity, the complement of a head-independent set is essential.

Proposition 4.1. *Suppose that the discrete time regulatory network $([0, 1]^d, F)$ satisfies coordinatewise injectivity. Let $(V, \mathcal{A})$ be the underlying network, let $U \subset V$ be head-independent, and let $W := V \setminus U$. Then for each $\mathbf{I} \in \mathcal{P}$, each $t \in \mathbb{N}$, and every (W, t) -specification $\mathbf{S}$ we have*

$$\#\{\mathbf{J} \in C(\mathbf{S}) : \mathbf{J} \subset \mathbf{I}\} \leq 1.$$

Proof. The statement holds for $t = 1$ since in this case if $\mathbf{J} \in \mathcal{Q}^1 \equiv \mathcal{P}$ is such $\mathbf{J} \subset \mathbf{I}$, then necessarily $\mathbf{J} = \mathbf{I}$. Now, given a $(W, 1)$ -specification $\mathbf{S}$ there are two possibilities: 1) either $\Pi_i \mathbf{S} = \Pi_i \mathbf{I}$ for each $i \in W$, or 2) there exists $i \in W$ such that $\Pi_i \mathbf{S} \cap \Pi_i \mathbf{I} = \emptyset$. In the first case $\#\{\mathbf{J} \in C(\mathbf{S}) : \mathbf{J} \subset \mathbf{I}\} = 1$, while in the second $\#\{\mathbf{J} \in C(\mathbf{S}) : \mathbf{J} \subset \mathbf{I}\} = 0$.

Let us suppose that $\#\{\mathbf{J} \in C(\mathbf{R}) : \mathbf{J} \subset \mathbf{I}\} \leq 1$ for $\tau \in \mathbb{N}$, for each (W, τ) -specification $\mathbf{R}$, and for every atom $\mathbf{I} \in \mathcal{P}$. Fix a $(W, \tau + 1)$ -specification $\mathbf{S}$. According to Lemma 3.3, if $\mathbf{J} \in \mathcal{Q}^{\tau+1}$ satisfies $\mathbf{S}$ then $P(\mathbf{J})$ satisfies $P(\mathbf{S})$. Therefore

$$\begin{aligned} \{\mathbf{J} \in C(\mathbf{S}) : \mathbf{J} \subset \mathbf{I}\} &\subset \{F(\mathbf{J}') \cap \mathbf{I} : \mathbf{J}' \in C(P(\mathbf{S}))\} \\ &= \cup_{\mathbf{I}' \in \mathcal{P}} \{F(\mathbf{J}') \cap \mathbf{I} : \mathbf{J}' \in C(P(\mathbf{S})) \text{ and } \mathbf{J}' \subset \mathbf{I}'\}. \end{aligned}$$

By hypothesis $\#\{F(\mathbf{J}') \cap \mathbf{I} : \mathbf{J}' \in C(P(\mathbf{S})) \text{ and } \mathbf{J}' \subset \mathbf{I}'\} \leq 1$ for each $\mathbf{I}' \in \mathcal{P}$. Now, for $\mathbf{I}' \in \mathcal{P}$ to be such that $\{F(\mathbf{J}') \cap \mathbf{I} : \mathbf{J}' \in C(P(\mathbf{S})) \text{ and } \mathbf{J}' \subset \mathbf{I}'\} \neq \emptyset$, it is needed that $\Pi_j \mathbf{I}' \supset \Pi_j P(\mathbf{S})$ for each $j \in W$, which already determines $\Pi_j \mathbf{I}'$ for $j \in W$. At the same time, for $\mathbf{J}' \subset \mathbf{I}'$ and $j \in W$ we have $\Pi_j(F(\mathbf{J}') \cap \mathbf{I}) = \mathbf{S}_j$. Therefore the system of constraints

$$(20) \quad \left(a\Pi_j \mathbf{J}' + (1-a) \sum_{i=1}^d K_{i,j} H(s_{i,j}(\mathbf{x}_i - T_{i,j})) \right) \cap \Pi_j \mathbf{I} = \Pi_j \mathbf{S}, \quad \forall j \in W,$$

must hold for any $\mathbf{x} \in \mathbf{J}'$. We consider this system of constraints as a system of equations for the unknown binary values $\{b_{i,j} := H(s_{i,j}(\mathbf{x}_i - T_{i,j})) : i \in U, (i,j) \in \mathcal{A}\}$. Since U is head-independent, each one of those equations contains at most one of these unknowns, and each unknown appears in at least one of those equations. Coordinatewise injectivity implies that for $j \in W$ fixed, the two possible values for $y_{i,j}$ appearing in (20) determine two disjoint intervals. Hence, if $b_{i,j} = b$ is a solution for (20), then $b_{i,j} = 1 - b$ is not. Thus, the system of constraints in (20) cannot be satisfied by two different choices $\{b_{i,j} \in \{0, 1\} : i \in U, (i,j) \in \mathcal{A}\}$. Finally, since each projection $\Pi_i \mathbf{I}'$ is completely determined by the values $\{H(s_{i,j}(\mathbf{x}_i - T_{i,j})) : (i,j) \in \mathcal{A}, \mathbf{x}_i \in \Pi_i \mathbf{J}'\}$, then the projections $\Pi_i \mathbf{J}'$, $i \in U$ are determined as well. In this way we fix the remaining projection $\Pi_i \mathbf{I}' \supset \Pi_i \mathbf{J}'$, proving that there is at most one atom $\mathbf{I}' \in \mathcal{P}$ such that $\#\{\mathbf{J}' \in C(P(\mathbf{S})) : \mathbf{J}' \subset \mathbf{I}', F\mathbf{J}' \cap \mathbf{I} \neq \emptyset\} = 1$. $\square$

Corollary 4.1 (Complement of a head-independent is essential). *Under the same hypotheses as in the previous proposition, the set $W := V \setminus U$ is essential.*

Proof. Indeed, since for each $I \subset \mathcal{P}$, each $t \in \mathbb{N}$, and every (W, t) -specification $\mathbf{S}$ we have $\#\{\mathbf{J} \in C(\mathbf{S}) : \mathbf{J} \subset \mathbf{I}\} \leq 1$, then $N(\mathbf{S}) = \sum_{\mathbf{I} \in \mathcal{P}} \#\{\mathbf{J} \in C(\mathbf{S}) : \mathbf{J} \subset \mathbf{I}\} \leq \#\mathcal{P}$, which proves the claimed result. $\square$

Remark 4.1. *The previous corollary holds if instead of the iterates of a coordinatewise injective map we consider a $\mathcal{P}$ -compatible sequence of $\mathcal{D}_{K,T,s,a}$, such that $U \subset V$ is head-independent with respect to the underlying network (V, A) associated to K .*

4.4. 2-loops. In order to have an effective reduction of degree in the polynomial bound for the complexity, we need that the underlying network admits redundant sets with essential complement. In general, head-independent sets are not redundant, and it seems more difficult to find sufficient conditions for vertices to be redundant than sufficient conditions for vertex sets to be essential. Below we will show a case where the structure of a two-vertices subnetwork is such that one of its vertices drives the other.

We say that the interaction matrix $K \in \mathcal{M}_{d \times d}([0, 1])$ is *non-degenerated* if for each $j \in \{1, 2, \dots, d\}$, the map

$$\left\{ \mathbf{e} \in \{0, 1\}^d : \mathbf{e}_i = 0 \text{ for all } i \text{ such that } K_{i,j} = 0 \right\} \ni \mathbf{e} \mapsto \sum_{i=1}^d \mathbf{e}_i K_{i,j} \in [0, 1]$$

is one-to-one.

Fix a network $(V, \mathcal{A})$. A *2-loop* in $(V, \mathcal{A})$ is a couple $\{i, j\} \subset V$ such that $(i, j) \in \mathcal{A}$ and i is the only tail for an arrow whose head is j , and $(j, i) \in \mathcal{A}$ and i is the only head for an arrow whose tail is j . In this case we say that j is the *isolated end of the 2-loop*.

Proposition 4.2. *Suppose that $([0, 1]^d, F)$ satisfies coordinatewise injectivity, and suppose that $\{i, j\} \subset V$ is a 2-loop in the underlying network $(V, \mathcal{A})$ whose isolated end is the vertex j . If in addition the interaction matrix K is non-degenerated, then i drives j .*

Proof. We will prove that, for each $t \in \mathbb{N}$, each $J \in \mathcal{Q}_i^t$, and every $I \in \mathcal{P}_j$, there exists at most one interval $J' \in \mathcal{Q}_j^t$ such that $J' \subset I$, and $C(J \times J') \neq \emptyset$. In this way we prove that i drives j with multiplicity $m_{i,j} \leq \#\mathcal{P}_j = 2$.

For $t = 1$ the statement obviously holds. Suppose it holds for $t = \tau$. Fix $J \in \mathcal{Q}_i^{\tau+1}$ and $I \subset \mathcal{P}_j$. Then, thanks to the coordinatewise injectivity, there exists a unique contraction $f \in \mathcal{F}_i$ and a unique interval $P(J) \in \mathcal{Q}_j^\tau$ such that $J = f(P(J)) \cap I'$. Here I' is the atom $\mathcal{P}_i$ which contains J .

The affine contraction $f : [0, 1] \rightarrow [0, 1]$ has the form $f(x) = ax + (1 - a) \sum_{k=1}^d \mathbf{e}_k K_{k,i}$, for some $\mathbf{e} \in \{0, 1\}^d$. Since the interaction matrix is non-degenerated, then the values of $\mathbf{e}_k \in \{0, 1\}$, for $K_{k,i} > 0$, are uniquely determined by f . Therefore, given $J \in \mathcal{Q}_i^{\tau+1}$, if $\mathbf{J}' = P(\mathbf{J})$ for some $\mathbf{J} \in C(J)$, not only we have $\mathbf{J}' \in C(P(J))$ as established in Lemma 3.3, but also $H(s_{k,i}(\mathbf{x}_k - T_{k,i})) = \mathbf{e}_k$, for each $\mathbf{x} \in \mathbf{J}'$.

From here on we use the fact that $\{i, j\}$ is a 2-loop with isolated end j , i. e., (a) $K_{j,k} = 0$ for $k \neq i$, and (b) $K_{k,j} = 0$ for $k \neq i$. Condition (a) implies that the value $\mathbf{e}_j \in \{0, 1\}$ determines the atom $I'' \in \mathcal{P}_j$ containing $\Pi_j \mathbf{J}'$ for each $\mathbf{J}' \in C(P(J))$. Here we apply the induction hypothesis. Given $P(J) \in \mathcal{Q}_j^\tau$ and $I'' \in \mathcal{P}_j$, there is at most one interval in $\mathcal{Q}_j^\tau$, which we denote J' , such that $J' \subset I''$ and $C(P(J) \times J') \neq \emptyset$. By condition (b), the image of each $\mathbf{J}' \in C(P(J) \times J')$ is the interval $\Pi_j F(\mathbf{J}') = aJ' + (1 - a)H(s_{i,j}(\mathbf{x}_i - T_{i,j}))$, which depends only on J and $P(J) \ni \mathbf{x}_i$. Thus, the interval $\Pi_j F(\mathbf{J}') \cap I \in \mathcal{Q}_j^{\tau+1}$ is uniquely determined from $J \in \mathcal{Q}_i^{\tau+1}$ and $I \in \mathcal{P}_j$. $\square$

Remark 4.2. *Once again, proposition 4.2 holds for compatible sequence of maps with common base partition, all of the satisfying piecewise injectivity, and having the same underlying network.*

5. EXAMPLES

5.1. Circuits. The circuit on d vertices has vertex set $V = \{1, 2, \dots, d\}$ and arrow set $\mathcal{A} \subset V \times V$ satisfying

$$(i, j) \in \mathcal{A} \Leftrightarrow j = i + 1, \text{ or } i = d \text{ and } j = 1.$$

Some numerical experiments suggest that in large random networks, and for a large class of initial conditions, the state of most of the units in the network converges to a fixed value, whereas in a small subnetwork, an oscillatory regime takes place. Very often this subnetwork is a circuit of low dimension. The circuit on d vertices may be negative or positive according to the value of the

product $\sigma := s_{d,1} \times \prod_{i=1}^{d-1} s_{i,i+1}$, defined by the activation matrix. In discrete time regulatory networks defined in positive circuits, numerical experiments have shown multistability, i. e., coexistence of multiple attractive fixed points. On the contrary, for the systems defined on negative circuits, beside some degenerated cases, the dynamics does not admit fixed points. A rigorous study of this numerically observed phenomena is in progress. Concerning the complexity of this kind of systems we can already state the following.




Figure 1.

A discrete time regulatory network whose underlying network is a circuit on d vertices, and whose contraction rate is smaller than $1/2$, satisfies $C(t) = \mathcal{O}(t^d)$. This follows from Theorem 3.1, taking into account that these systems satisfy coordinatewise injectivity. Indeed, since the interaction matrix is normalized, i. e., $\sum_{i=1}^d K_{i,j} = 1$, then $K_{i,j} = 1$ for each $(i, j) \in \mathcal{A}$. Using the same notation as in Proposition 3.1, we have that in this case $\mathcal{I}_i = \{0, 1\}$ for each $i \in \{1, 2, \dots, d\}$, implying that $a_0 = 1/2$.

The genetic toggle switch described in [12] is a concrete experimental example of a positive 2-circuit, made of two mutually inhibiting interacting units. In this case each unit is composed by a couple promoter–repressor, such that each promoter is inhibited by the repressor transcribed by the opposite promoter. This system is traditionally modeled by the system of ordinary differential equations

$$\begin{aligned} \frac{dr_1}{dt} &= \frac{\alpha_1}{1 + r_2^{\beta_2}} - r_1 \\ \frac{dr_2}{dt} &= \frac{\alpha_2}{1 + r_1^{\beta_1}} - r_2, \end{aligned}$$


where r_i represents the concentration of repressor i , α_i is the effective rate of synthesis of repressor i , and β_i is the cooperativity of repression of promoter i . The corresponding network of piecewise

contractions has the form

$$\begin{aligned} r_1(t+1) &= ar_1(t) + (1-a)H(T_{1,2} - r_2(t)) \\ r_2(t+1) &= ar_2(t) + (1-a)H(T_{2,1} - r_1(t)), \end{aligned}$$

where the rate of synthesis of the repressors is controlled by the parameter a , and the cooperativity of repression of promoters is related to both a and the thresholds $T_{i,j}$. The dynamical behavior of the positive circuit in two vertices is completely understood (see [3]), and it can be either bistable (the same as the continuous model) or conjugated to a rotation. In all cases the dynamical complexity asymptotically bounded by $t \mapsto t + 1$.

For the general circuit on 2 vertices, if the contraction rate is smaller than $1/2$, each one of the 2 vertices is at the same time redundant and essential. Indeed, in this case the coordinatewise injectivity holds, each vertex constitutes a head-independent set, and at the same time it is the isolated end of a 2-loop. Corollary 4.1 and Proposition 4.2 apply, implying that each vertex is redundant and essential at the same time. Hence, a system of two interacting units whose underlying network is a circuit on 2 vertices, and whose contraction rate is smaller than $1/2$, necessarily has linear complexity. Furthermore, for the negative circuit, since the image of rectangle in $\mathcal{Q}^\tau$ cannot intersect the double discontinuity ($\mathbf{x}_1 = T_{1,2}, \mathbf{x}_2 = T_{2,1}$), then it cannot have more than one successor. Taking into account this constraint in the computations we performed above, we deduce that in this case $C(t) \leq 2t + 2$ for each $t \in \mathbb{N}$. It has been proved [3] that the negative circuit admits an arbitrary large number of periodic orbits, as one moves on the line $\{(T, a) : T_{1,2} = T_{2,1} = 1/2, a \in [1/2, 1)\}$. There the number of admissible periodic orbits grow as one increases the contraction rate. Because of this, the complexity increases as a grows, though our numerical experiments indicate that $C(t)$ can always be bounded by a quadratic polynomial, as it is illustrated in Figure 2. We conjecture that the complexity of the negative circuit on 2 vertices can always be bounded by a linear function $C(t) \leq K \times (t + 1)$, where the constant factor depends on the choice of parameters (T, a) , and it diverges as we increase a .

**Figure 2.**

Log-log plot of the complexity of the negative circuit on 2 vertices for $T_{1,2} = T_{2,1} = 0.5$ and $a = 0.93$. The straight line is the log-log plot of the quadratic upper bound $t \rightarrow 4t^2$.

For a circuit on d vertices any independent set of vertices (i. e., no two of them are adjacent to the same arrow) is head-independent. In particular $\{2, 4, \dots, 2\lfloor d/2 \rfloor\}$ is a maximal independent set, therefore, according to Corollary 4.1 its complement $\{1, 3, \dots, 2\lfloor (d-1)/2 \rfloor + 1\}$ is essential. If we think of the collection $\mathcal{Q}^\tau$ as a finite time version of the attractor, Proposition 4.1 establishes that this coarse grained attractor has dimension $k = \lfloor d/2 \rfloor$. Indeed, in this case each d -dimensional interval of $\mathbf{J} \in \mathcal{Q}^\tau$ has at most k independent one-dimensional projections, whereas $d-k$ projections are determined by the first, and by the atom of $\mathcal{P}$ containing $\mathbf{J}$. In this way, the coarse grained version of the attractor lies inside the graph of a $\#\mathcal{P}$ -valued function defined on $[0, 1]^k$, and taking values on $[0, 1]^{d-k}$. Note that this property holds uniformly on τ , suggesting that the attractor itself may be embedded in the graph of a multivalued function. This of course has yet to be proved.

Let us remark that in [9] it is presented an biological example of a negative circuit on 3 vertices, the so-called repressiliator. They report the existence of oscillatory behavior in time, which would correspond to periodic solutions of the corresponding model, though the period and amplitude exhibit significant variability. This may correspond either to stochastic effects, or may reflect intrinsic complex behavior.

5.2. Networks with loops. With respect to the effective reduction of the polynomial upper bound, the example of the circuit on 2 vertices with contraction rate smaller than $1/2$ may be generalized as follows. Fix a network $(V, \mathcal{A})$ and 2-loops $\{i, j\}$ and $\{i', j'\}$, with isolated ends j and j' respectively. We say that “ $\{i, j\}$ and $\{i', j'\}$ are disjoint” if $\{k \in V : (i, k) \in \mathcal{A}\} \cap \{k \in V : (i', k) \in \mathcal{A}\} = \emptyset$, i. e., the non-isolated ends have disjoint heads.

Let $(F, [0, 1]^d)$ be a network with non-degenerated interaction matrix (as defined in the previous section), and satisfying the coordinatewise injectivity. If the underlying network $(V, \mathcal{A})$ has $k := d$ pairwise disjoint 2-loops, then $C(t) = \mathcal{O}(t^{d-k})$. Indeed, let $\{\{i_n, j_n\} : n = 1, 2, \dots, k\}$ be the set of mutually disjoint 2-loops, and let j_n is the isolated end of $\{i_n, j_n\}$ for each $n = 1, 2, \dots, k$. According to Proposition 4.2, the set $U := \{i_n : n = 1, 2, \dots, k\}$ is redundant. Indeed, each j_n drives i_n for each $n = 1, 2, \dots, k$. On the other hand, since the 2-loops are pairwise disjoint, the set U head-independent too, and so by Corollary 4.1 the complement $W := V \setminus U$ is essential, therefore $C(t) = \mathcal{O}(t^{d-k})$.





Figure 3.

In Figure 3 we represent a network containing 3 disjoint 2-loops with isolated ends a , b and c . According to Theorem 3.1 and Corollary 4.2, if the contraction rate is sufficiently small and the interaction matrix is non-degenerated, then the complexity of the system is bounded by a cubic polynomial.

In [21] a discrete time regulatory network of the type considered in this paper was used to model the regulation of the expression of the tumor suppressor gene *p53*, whose product acts as an inhibitor of uncontrolled cell growth. The underlying network of this model is shown in Figure 4. The node m represents the expression level of the *mdm2* gene, whose product inhibits the expression of *p53*. Nodes b and c represent blood-vessel formation and cell proliferation respectively. Though b and c are not expression levels of a particular gene, they represent relevant quantities directly implicated in cancer disease.

**Figure 4.**

Regulatory network for the *p53* gene expression. The arrow indicates positive interaction, while $\dashv$ indicates an inhibitory interaction.

In this network the couple *p53*–*m* form a 2-loop with isolated node *m*, and the vertices *m* and *c* are head-independent. If we assume coordinatewise injectivity, according to proposition 4.1, nodes *p53* and *c* capture the whole complexity of the network. On the other hand, since node *m* is redundant, the complexity of the network is bounded by a polynomial of degree 3. Note that in this case the network admits the base-bundle decomposition $\{p53, m\} \cup \{b, c\}$, and both base ($V_b := \{p53, m\}$) and bundle ($V_d := \{b, c\}$) define circuit on two vertices. Therefore, if coordinatewise injectivity holds, Theorem 4.1 implies that the complexity of this four-vertices network is bounded by a quadratic polynomial.

6. FINAL REMARKS

In this work we formulate and show the first results towards the solution of the following theoretical problem: determine to what extent the topological structure of the underlying network determines the behavior of a discrete time regulatory network. By solving this problem we intend to acquire a better understanding of the behavior observed in real life regulatory networks. The next step in our program consist in extending Theorem 3.1 to the case of arbitrary contraction rates, when coordinatewise injectivity does not hold. Some preliminary numerical experiments support lead us to conjecture that the complexity is always bounded by a polynomial; furthermore, this experiments suggest that the asymptotic dynamics consist of a finite number of periodic or quasiperiodic orbits. If this is true, the complexity should be always bounded by a polynomial of degree one.

Though in general it is not possible to deduce relevant dynamical properties only from the complexity, we expect that in our case the growth of the complexity could give information about topological and recurrence properties of the attractor. For instances, the growth rate of the complexity could be related to the number of transitive components of the attractor. Anyway, in order

to have a complete description of the qualitative behavior of our models we need to consider other characteristics besides the dynamical complexity.

Theorems 4.2 and 4.1 we have specific instances of constraints imposed on the behaviors of the system by the structure of the underlying network. In this direction, our next goal is to study how the behavior of a network is related to the possible dynamical behaviors of its subnetworks. Results in this direction would allow us to deal with questions such as the role of self-regulation in the stabilization of the circuits, and in general, to progress in the understanding of the engineering behind the design of real life functional networks.

REFERENCES

- [1] A. Becskei and L. Serrano, Engineering stability in gene networks by autoregulation, *Nature* **405** (2000) 590–592.
- [2] F. Blanchard, B. Host, and A. Maass, Topological complexity, *Ergodic Theory and Dynamical Systems* **20** (2000) 641–662.
- [3] R. Coutinho, B. Fernandez, R. Lima and A. Meyroneinc, Discrete time piecewise affine model of genetic regulatory networks, Preprint CPT 2004.
- [4] R. Coutinho, Dinâmica Simbólica Linear, Ph.D. Thesis, Instituto Superior Técnico-Universidade Técnica de Lisboa 1999.
- [5] J. Cassaigne, Recurrence in infinite words, Proceedings of the 18th Symposium on Theoretical Aspects of Computer Science (STACS 2001), Dresden (Allemagne), *Lecture Notes in Computer Science* **2010** (2001) 1–11.
- [6] H. de Jong, Modeling and simulation of genetic regulatory systems: A literature review, *Journal of Computational Biology* **9**(1) (2002) 69–105.
- [7] H. de Jong and R. Lima, Modeling the dynamics of genetic regulatory networks: Discrete and continuous approaches, CML2004 Lecture Notes to appear.
- [8] R. Edwards, Analysis of continuous-time switching networks, *Physica D* **146** (2000) 165–199.
- [9] M. B. Elowitz and S. Leibler, A synthetic oscillatory network of transcriptional regulators *Nature* **403** (2000) 335–338.
- [10] S. Ferenczi, Complexity of sequences and dynamical systems, *Discrete Mathematics* **206** (1999) 145–154.
- [11] J.-M. Gambaudo and Ch. Tresser, On the dynamics of Quasi-contraction, *Bol. Soc. Bras. Mat.* **19** (1)(1988) 61–114.
- [12] T. S. Gardner, C. R. Cantor and J. J. Collins, Construction of a genetic toggle switch in *Escherichia coli*, *Nature* **403** (2000) 339–342.
- [13] L. Glass, Combinatorial and topological methods in nonlinear chemical kinetic, *J. Chem. Phys.* **63** (1975) 1325–1335.
- [14] B.C. Goodwin, “Temporal Organization in Cells”, *Academic Press*, 1963.
- [15] R. Hengge-Aronis, Signal transduction and regulatory mechanisms involved in control of the σ^S (RpoS) subunit of RNA polymerase, *Microbiology and Molecular Biology Reviews* **66** (3) (2002) 373–395.
- [16] T. Ideker and T. Galitski and L. Hood, A new approach to decoding life: Systems biology, *Annual Review of Genomics and Human Genetics* **2** (2001) 343–372.
- [17] H. Kitano, Systems biology: A brief overview, *Science* **295** (5560), (2002) 1662–1664.
- [18] D. Thieffry and R. Thomas, Dynamical behavior of biological networks, *Bulletin of Mathematical Biology* **57** (2)(1995) 277–297.
- [19] R. Thomas, Boolean formalization of genetic control circuits, *J. Theor. Biol.* **42** (1973) 563–585.
- [20] R. Thomas and M. Kaufman, Multistationarity, the basis of cell differentiation and memory II: Logical analysis of regulatory networks in terms of feedback circuits, *Chaos* **11** (1)(2001) 180–195.
- [21] Carlos Aguirre, João Martin and R. Vilela Mendes, Dynamics and coding of a biologically-motivated network, q-bio.MN/0407031 to appear in *Int. J. Bifurcation and Chaos*.